



Global Biogeochemical Cycles

RESEARCH ARTICLE

10.1029/2018GB005880

Key Points:

- New simple parameterization for production of bioavailable P from acid dissolution of mineral dust incorporated into a global aerosol model
- Inclusion of acid dissolution increases atmospheric flux of bioavailable P from dust to oceans by 120% and drives dust P bioavailability
- Major increases in bioavailable P from atmospheric acid processes occur in Mediterranean Sea, North Atlantic, NW Pacific, and Indian Ocean

Supporting Information:

- Supporting Information S1

Correspondence to:

R. J. Herbert,
r.j.herbert@reading.ac.uk

Citation:

Herbert, R. J., Krom, M. D., Carslaw, K. S., Stockdale, A., Mortimer, R. J. G., Benning, L. G., et al. (2018). The effect of atmospheric acid processing on the global deposition of bioavailable phosphorus from dust. *Global Biogeochemical Cycles*, 32, 1367–1385. <https://doi.org/10.1029/2018GB005880>

Received 7 JAN 2018

Accepted 9 AUG 2018

Accepted article online 20 AUG 2018

Published online 22 SEP 2018

This article was corrected on 2 OCT 2018. See the end of the full text for details.

The Effect of Atmospheric Acid Processing on the Global Deposition of Bioavailable Phosphorus From Dust

R. J. Herbert^{1,2} , M. D. Krom^{1,3}, K. S. Carslaw¹ , A. Stockdale¹ , R. J. G. Mortimer⁴ , L. G. Benning^{1,5,6} , K. Pringle¹, and J. Browse^{1,7}

¹School of Earth and Environment, University of Leeds, Leeds, UK, ²Now at Department of Meteorology, University of Reading, Reading, UK, ³Now at Morris Kahn Marine Research Station, Department of Marine Biology, Leon H. Charney School of Marine Science, University of Haifa, Haifa, Israel, ⁴School of Animal, Rural and Environmental Sciences, Nottingham Trent University, Nottinghamshire, UK, ⁵GFZ German Research Centre for Geosciences, Potsdam, Germany, ⁶Department of Earth Sciences, Free University of Berlin, Berlin, Germany, ⁷Now at College of Life and Environmental Sciences, University of Exeter, Penryn, UK

Abstract The role of dust as a source of bioavailable phosphorus (Bio-P) is quantified using a new parameterization for apatite dissolution in combination with global soil data maps and a global aerosol transport model. Mineral dust provides 31.2 Gg-P/year of Bio-P to the oceans, with 14.3 Gg-P/year from labile P present in the dust, and an additional 16.9 Gg-P/year from acid dissolution of apatite in the atmosphere, representing an increase of 120%. The North Atlantic, northwest Pacific, and Mediterranean Sea are identified as important sites of Bio-P deposition from mineral dust. The acid dissolution process increases the fraction of total-P that is bioavailable from ~10% globally from the labile pool to 18% in the Atlantic Ocean, 42% in the Pacific Ocean, and 20% in the Indian Ocean, with an ocean global mean value of 22%. Strong seasonal variations, especially in the North Pacific, northwest Atlantic, and Indian Ocean, are driven by large-scale meteorology and pollution sources from industrial and biomass-burning regions. Globally constant values of total-P content and bioavailable fraction used previously do not capture the simulated variability. We find particular sensitivity to the representation of particle-to-particle variability of apatite, which supplies Bio-P through acid-dissolution, and calcium carbonate, which helps to buffer the dissolution process. A modest 10% external mixing results in an increase of Bio-P deposition by 18%. The total Bio-P calculated here (31.2 Gg-P/year) represents a minimum compared to previous estimates due to the relatively low total-P in the global soil map used.

Plain Language Summary Phosphorus (P) is an essential requirement for life. Natural sources of P on land are from rock weathering and fertilizers. By contrast over the open ocean, the major source of P is from falling dust. However, less than 10% of the P in dust is automatically available to phytoplankton for growth, a percentage we call bioavailable-P. Therefore, changes to the supply of bioavailable-P to oceans can have considerable impacts on marine ecosystems and the global carbon cycle. Previous work shows acid processes in the atmosphere can convert nonbioavailable minerals to bioavailable-P. In our previous study we found a simple relationship between acid in the atmosphere and bioavailable-P formed. Here we use this new relationship, together with global soil data maps on the amount and type of P in dust and a global aerosol transport model, which predicts where dust and acid interact. We calculate how much and where acid-modified dust ends up in the ocean. We show atmospheric acid processing of dust is particularly important in the Mediterranean Sea, North Atlantic Ocean, northwest Pacific Ocean, and the Indian Ocean. As a result, atmospheric acid pollution increases the amount of oceanic plant growth and reduces the quantity of atmospheric anthropogenic carbon dioxide.

1. Introduction

Phosphorus (P) is an essential requirement for life. In the terrestrial ecosystem P is made accessible naturally through weathering processes and increasingly through the supply of fertilizer and other P-containing compounds from anthropogenic processes. The dominant supply of P to coastal systems is from rivers and wastewater discharge, which are augmented by sedimentary recycling processes (Ruttenberg, 2003). By contrast, in the offshore ocean the dominant external supply is from atmospheric deposition processes, with long-range transportation of mineral dust from desert regions being the most important single source of P to the ocean surface (Graham & Duce, 1982; Mahowald et al., 2008; Myriokefalitakis et al., 2016). In situ measurements of

atmospheric aerosol (e.g., Carbo et al., 2005; Herut et al., 1999; Zamora et al., 2013) and soil mineralogical databases (Nickovic et al., 2012; Yang et al., 2013) show that total P (hereafter TP) content of soils and mineral dust is spatially and temporally variable and may range from <500 to >1,000 ppm, with corresponding variability in the component P pools.

Many studies show the importance of atmospheric supply in general and dust in particular as a source of external nutrients (Fe, N, and P) to the open ocean (Krishnamurthy et al., 2010; Mahowald et al., 2008). In such systems, the total N supplied is in a readily bioavailable form. In contrast, the P (and Fe) is supplied in both labile (and hence bioavailable) and nonbioavailable forms. The nonbioavailable particles of P can drop through the photic zone without enhancing phytoplankton growth and hence carbon uptake. Herut et al. (2005) showed that when fresh Saharan dust and dust pretreated to remove any water-leachable nutrients were added to a microcosm experiment containing N- and P-limited Mediterranean seawater only the fresh dust resulted in increased chlorophyll content. The fresh dust added nutrients in the ratio of 31N:1P (nitrate:phosphate), and thus, it was the water-leachable fraction of the dust rather than the particle itself that caused the extra productivity in this N and P colimited system (Thingstad et al., 2005). Eijssink et al. (2000) found that even in the P-limited eastern Mediterranean, 70% of the TP (mainly detrital apatite) supplied as Saharan dust to the surface waters was transferred through the water column and ended up deposited in the sediment. The remaining 30% was taken up by biological processes in the water column.

Observations suggest that the percentage of TP in mineral dust that is deposited in a bioavailable form (Bio-P) is spatially variable, ranging from <10 to >80% (Baker, French, et al., 2006; Markaki et al., 2003; Vet et al., 2014; Zamora et al., 2013), and may increase with distance from the dust source (Baker, French, et al., 2006). Recent ambient observations and laboratory experiments (Nenes et al., 2011; Srinivas & Sarin, 2015; Stockdale et al., 2016) provide support for the hypothesis that atmospheric acidification and subsequent dissolution are a primary process for producing Bio-P in mineral dust. Increasing the amount of leachable, or bioavailable, P by acid processes in the atmosphere will have a direct effect of increasing phytoplankton biomass and hence carbon uptake in a variety of ocean systems (e.g., Jickells & Moore, 2015; Mahowald et al., 2008).

The paucity of observational data from the open oceans necessitates the use of models to determine the atmospheric flux of Bio-P to surface waters. Models are also needed to quantify how Bio-P deposition may change with environmental factors such as source strength, atmospheric pollution, and large-scale changes to global circulation. To achieve this, models must represent the emission, transport, and deposition of the particulate P, as well as processes that determine the TP and Bio-P content. Previous global deposition modeling studies using global chemical transport models (Brahney et al., 2015; Krishnamurthy et al., 2010; Mahowald et al., 2008; Wang et al., 2015) assumed that dust emissions contain a constant fraction of TP (between ~720 and 1,050 ppm) and a constant percentage of TP deemed Bio-P (between 10 and 15%). The result is a spatially varying flux of dust-borne P to the surface with a constant solubility, which may not reflect underlying variability between dust sources, nor atmospheric acidification processes. A recent study by Myriokefalitakis et al. (2016) represented the acidification process using an acid-solubilization mechanism in which apatite dissolution is treated as a kinetic process dependent on the H⁺ activity of the surrounding water droplet and known acid dissolution thermodynamic constants. Their study used a global soil mineralogy data set (Nickovic et al., 2012) to represent the spatially varying soil TP content with dust-TP emissions adjusted to a global mean of 880 ppm and a soluble fraction of 10% representing a leachable inorganic pool of P in the initial dust. The remaining TP was assumed to be apatite. This treatment reproduces the atmospheric acidification process but relies on several assumptions for estimating the different forms of P. A recent global soil database (Yang et al., 2013) provides high-resolution information on the geographical distribution of TP in its different forms including apatite, labile, organic, occluded, and secondary P. The data set shows that there is considerable spatial variability in all components. This data set provides more appropriate forms of P for use in estimating dust-borne P emissions and simulating atmospheric acidification processes of apatite.

A recent study by Stockdale et al. (2016) presents results from a series of experiments simulating atmospheric acidification on dust samples and dust precursor soils in which the acidity of the solution and the mass of dust was systematically varied. Phosphorus speciation experiments identified the dominant forms of phosphorus in the dust samples as ~80% apatite P, ~10% Fe-bound P, and ~10% labile P. The acidification experiments provide robust evidence for a relatively simple relationship in which the dissolution behavior of

mineral dust is controlled by the absolute number of protons in the solution, the calcium carbonate (CaCO_3) content, and the apatite (Ap-P) content of the dust. Rapid acid dissolution of both mineral species occurs simultaneously when both minerals are present on the same mineral grain (internally mixed). The dissolution rate of Ap-P is greater when the Ap-P and CaCO_3 exist on different grains (externally mixed). As a result of slow mineral precipitation kinetics the dissolved phosphate remains in solution when the H^+ concentration falls or water content increases. The total mass of Bio-P upon deposition to the surface is thus the sum of acid-dissolved apatite (Acid-P) and the loosely bound labile P (Lab-P) that was initially available in the particles. As the dissolution of Ap-P occurs at a faster rate when CaCO_3 is not present, if Ap-P and CaCO_3 were exclusively present on different particles (i.e., externally mixed), then it is possible that more Acid-P would be produced. As discussed by Stockdale et al. (2016) it is hypothesized that many particles would contain both Ap-P and CaCO_3 ; however, there is likely a degree of variation between particles with some containing more or less of each mineral component.

It is widely recognized that mineral dust plays an important role in the transport of nutrients to the open ocean surfaces; however, as shown by Yang et al. (2013) there exists considerable variability in the relative abundance of different P-containing components from different source regions. The results of Stockdale et al. (2016) additionally show that the spatial distribution of CaCO_3 and its abundance in relation to Ap-P may have considerable impacts on the production of Bio-P from acid dissolution of Ap-P. The simplistic dissolution mechanism presented by Stockdale et al. (2016) and the detailed soil-P speciation data set presented by Yang et al. (2013) provide a new and readily applicable method for estimating the spatiotemporal distribution of Bio-P from dust and allows us to investigate the sensitivity of Acid-P production to the degree of internal and external mixing of the components. This will help to understand key uncertainties in the acid dissolution process and help better define the focus of future research.

In this study we use the Global Model of Aerosol Processes (GLOMAP) coupled to the global chemical transport model TOMCAT to simulate the emission, transport, and deposition of dust-borne phosphorus to the surface. Apatite dissolution parameterizations based on the results from Stockdale et al. (2016) and a database of soil P speciation (Yang et al., 2013) are used to simulate atmospheric acidification of mineral dust by H_2SO_4 and HNO_3 ; the soil database is also used to simulate the contribution of the dust-borne Lab-P as a spatially variable percentage of dust mass. These results are used to investigate the drivers of the spatiotemporal variability in dust-borne Bio-P and the Acid-P (the amount of TP converted to Bio-P by atmospheric acid processes). The results are compared to a global data set of observations (Vet et al., 2014) and results from other modeling studies to help understand the importance of atmospheric acidification of mineral dust on the global supply of P nutrients to the open oceans and elsewhere. Finally, simulations are performed to investigate the sensitivity of the results to the effect of external mixing of apatite and CaCO_3 on the amount of Bio-P delivered to the ocean surface.

2. Model Description

We use the global aerosol microphysics model GLOMAP-mode (Mann et al., 2010) coupled to the 3-D global chemical transport model TOMCAT (Chipperfield, 2006). Simulations are performed using a horizontal resolution of 2.8° by 2.8° and 31 hybrid σ -pressure levels extending from the surface to 10 hPa. Meteorology for the simulated year is driven by the European Centre for Medium-Range Weather Forecasts (ECMWF) ERA-Interim reanalyses at 6-hourly intervals. Monthly mean low-cloud fields are prescribed from the International Satellite Cloud Climatology Project (ISCCP) archive. In the standard GLOMAP-mode setup the aerosol particle number and size distribution are described using a modal scheme with seven internally mixed modes (four soluble and three insoluble) and five aerosol species: sulfate, black carbon, organic carbon, sea salt, and dust. The seven modes cover four size ranges: nucleation (~ 1 to 10 nm diameter), Aitken (~ 10 to 100 nm), accumulation (~ 0.1 to 0.5 μm), and coarse (~ 0.5 to 10 μm).

The aerosol processes treated in the model include primary and precursor emissions, nucleation of H_2SO_4 aerosol, sedimentation and dry deposition, hygroscopic growth, in-cloud activation and scavenging, below-cloud scavenging, intermodal and intramodal coagulation, condensation of H_2SO_4 and secondary organic vapors onto all aerosol particles, aging of water-insoluble particles, and in-cloud processing (formation of sulfate mass from oxidation of sulfur dioxide in aerosol particles activated into cloud droplets).

GLOMAP-mode also includes an aerosol precursor chemistry scheme primarily for simulating gas and aqueous phase reactions leading to the production of SO_2 and H_2SO_4 .

Monthly emissions of anthropogenic BC, OC, and SO_2 , and volcanic SO_2 are supplied by global data sets from AeroCom hindcast (Diehl et al., 2012). The Global Fire Emissions Database (GFED) v2 database (van der Werf et al., 2010) is used for wildfire emissions, and dimethyl sulfide (DMS) emissions are calculated using DMS seawater concentrations from Kettle et al. (1999) and transfer velocity parameterizations of Nightingale et al. (2000). Daily size-resolved emissions of dust (covering accumulation and coarse modes) are included via prescribed fluxes from AeroCom (Dentener et al., 2006). Deposition of aerosol species occurs via dry deposition, using a dry deposition velocity following Slinn (1982), and wet deposition through nucleation scavenging from large-scale and convective precipitation (activation of cloud condensation nuclei), impaction scavenging by precipitation, and scavenging by low-cloud drizzle (Browse et al., 2012). The aqueous chemistry scheme includes the dissolution of SO_2 , H_2O_2 , and O_3 into cloud droplets and the subsequent oxidation of S (IV) to S (VI) by H_2O_2 and O_3 . Condensation of H_2SO_4 onto aerosols is simulated using coefficients following Fuchs and Sutugin (1971) with added correction factors for molecular effects and limiting interfacial mass transport, the latter using an accommodation coefficient 1.0 for both insoluble and soluble modes as per Mann et al. (2010). For this study, the condensation of HNO_3 onto dust has been added following the above method; HNO_3 concentrations are taken from the coupled TOMCAT model with condensation acting as a sink to the HNO_3 fields. For simplicity, the condensed HNO_3 is added to the sulfate component as an equivalent mass of H_2SO_4 that would produce the same mass of H^+ ions upon dissociation of HNO_3 .

In this study we are interested in the sulfate associated with dust aerosol, rather than the sulfate associated with sea-salt, and thus have altered the model setup to isolate the dust-associated sulfate, which will be used to determine the amount of associated acid. Full details can be found in Text S1. The resulting annual dust deposition flux, shown alongside the data from the standard setup, is compared to AeroCom observations (Huneeus et al., 2011) in Figure S1 and shows that the current setup is able to simulate dust deposition fluxes both close to source and in remote regions and is comparable to the original GLOMAP setup.

To simulate the acid dissolution process in GLOMAP, two soil databases were used to provide information on the dust mineralogy. For the different forms of P in the dust the soil phosphorus database presented by Yang et al. (2013) was used. The database uses a global distribution of soil parent material and lithological dependent P concentrations along with a pedogenesis transformation model to simulate the weathering of each parent rock type. The authors present global distributions of P in different forms as an average of the top 50 cm of the soil profile, including total P (TP), labile inorganic P (Lab-P), and apatite-P (Ap-P). A database of global soil CaCO_3 content was obtained from the gridded Global Soil Dataset for use in Earth System Models (GSDE) developed by Shangguan et al. (2014). A mean content was determined using the top 50 cm of data in order to provide comparable emissions to the P components and to maximize global coverage of emissions. Figure 1 shows the data sets regridded to a 2.8° horizontal resolution. Dust TP content typically ranges from 200 to 1,000 ppm by mass (global mean of 563 ppm), Ap-P from 20 to 300 ppm (164 ppm), and Lab-P from 20 to 100 ppm (50 ppm), while CaCO_3 content, at over two orders of magnitude greater, typically ranges from 1 to 12% (global mean of 3.3%) of the total dust mass. In terms of global means Lab-P constitutes 9% of TP dust content and varies between 5 and 10%, whereas Ap-P constitutes 29% of TP and varies between 5 and 60%; the major deserts show less variability and Ap-P tends to represent $\sim 50\%$ of TP. The form of P that represents the majority of remaining TP is occluded-P (not shown; spatial distribution apparent from the low values of Ap-P:TP). Experimental observations from Stockdale et al. (2016) suggest that this form of P is not released through acid dissolution nor upon deposition to the ocean surface; it is also much more prevalent in the northern boreal regions where dust uplift is less important.

To simulate the emission and deposition of the dust mineralogy, tracers were included in GLOMAP alongside dust following previous studies (Atkinson et al., 2013; Vergara-Temprado et al., 2017) using the regridded soil-P and CaCO_3 data sets. For each tracer the spatially variable mass fraction of the components from the regridded soil-P and CaCO_3 databases were applied to the modeled dust emission at each grid point to produce an associated mass. Throughout the simulation, dust is considered a single component with the fractional partition between the constituents conserved for all processes, including interaction between two dust modes. The simulations produce monthly mean wet and dry mass deposition fluxes (both impaction




Figure 1. Global gridded data sets used for Global Model of Aerosol Processes dust-associated emissions of (a) total-phosphorus (ppm), (b) apatite-phosphorus (ppm), (c) labile-phosphorus (ppm), (d) calcium-carbonate (%), (e) ratio of apatite-phosphorus to total-phosphorus (%), and (f) ratio of labile-phosphorus to total-phosphorus (%).

and nucleation scavenging) from each grid point for total dust, TP, Ap-P, Lab-P, CaCO₃, and dust-associated acid in the form of the sulfate component.

The acid dissolution process is simulated using parameterizations based on experimental observations from Stockdale et al. (2016), which provide robust evidence that the dissolution of Ap-P occurs rapidly (seconds to minutes) and that the system can be replicated by a simple linear relationship between acid input and Ap-P/CaCO₃ dissolution. Linear fits to the experimental data on a logarithmic scale using all dust samples provide the mass of dissolved Ap-P and CaCO₃ as a function of the absolute number of protons available. The dissolution rate of Ap-P increases in the absence of CaCO₃; therefore, an additional step has been introduced to simulate the enhanced dissolution rate. Stockdale et al. (2016) found that the dissolution of hydroxyapatite powder was well reproduced by the geochemical PHREEQC model (Parkhurst & Appelo, 2013); this model was used to produce a linear fit between available protons and dissolution of Ap-P, which provides a means for estimating Ap-P dissolution in the absence of CaCO₃. The resulting parameters from the linear fits can be found in Table S1. At each grid point (horizontal and vertical) the number of protons available, determined from the mass of dust-associated sulfate and assuming each mole of H₂SO₄ dissociates to produce 2 moles of H⁺, is used to determine the number of moles of deposited dissolved CaCO₃ (mol CaCO_{3diss}) using equation (1). Applying mol CaCO_{3diss} to equation (2) provides dissolved Ap-P (mol Ap-P_{diss}) from simultaneous dissolution of Ap-P and CaCO₃. If all CaCO₃ has been dissolved (i.e., mol CaCO_{3diss} = mol CaCO_{3available}), any excess protons are used to determine a second source of mol Ap-P_{diss} using equation (2) under the condition that mol CaCO_{3available} = 0. The number of moles of dissolved Ap-P from simultaneous and separate dissolution, constrained by the available deposited mass of Ap-P, is integrated over each horizontal grid point to provide a monthly mean surface deposition flux of bioavailable P from Ap-P dissolution. The deposition fluxes of Lab-P are similarly integrated over each grid point and added to the dissolved Ap-P to provide the total atmospheric bioavailable P flux to the surface from inorganic mineral dust.

$$\log_{10}(\text{mol CaCO}_{3\text{diss}}) = a_1 + b_1 \cdot \log_{10}(\text{mol H}^+) \quad (1)$$

$$\log_{10}(\text{mol Ap-P}_{\text{diss}}) = \begin{cases} a_2 + b_2 \cdot \log_{10}(\text{mol CaCO}_{3\text{diss}}), & \text{if mol CaCO}_{3\text{available}} > 0 \\ a_3 + b_3 \cdot \log_{10}(\text{mol H}^+_{\text{excess}}), & \text{if mol CaCO}_{3\text{available}} = 0 \end{cases} \quad (2)$$

3. Results

3.1. Bioavailable Phosphorus

Annual deposition fluxes of bioavailable phosphorus (Bio-P) from dust, dust mass, and the dust-associated acid for the year 2001 are shown in Figure 2. The large spatial variability in dust deposition drives the spatial variability of Bio-P. Highest values are found close to major dust source regions of Sahara and Kalahari deserts in Africa, and across central Asia, the Americas, and Australia. A significant proportion of the dust is deposited close to the source of emission. However, considerable transport occurs from all dust source regions driven by large-scale meteorology, most notably the transport of African dust advecting westward across the Atlantic Ocean driven by trade winds and midlatitude transport from South America and South Africa eastward. Northern Hemisphere midlatitude transport from Asia across the western and central Pacific Ocean also provides a considerable transport route for dust to these remote ocean regions. The amount of acid condensed onto the mineral dust (Figure 2c) shows a strong correlation with regions of high pollution and dust transport. The industrialized regions of North America, Europe, and central and eastern Asia can be clearly identified, as well as regions of biomass burning in central Africa. Transported pollutants also play a considerable role over the north-west Pacific and North Atlantic, as well as northern Africa and the Mediterranean Sea. The low dust loading in the central Pacific and Atlantic coincident with a moderate amount of acid highlights the marine source of sulfate emissions from DMS in this region, and a similar correlation in the northern high latitudes highlights the transport of pollutants from Europe and North America to this remote region.

Table 1 provides a breakdown of absolute mass and mass per unit area deposited per year globally, on terrestrial ecosystems, on marine ecosystems, and into each ocean basin and continent; boundaries to each basin are shown in Figure S2. Globally, 117 Gg-P/year Bio-P from mineral dust is deposited to the surface annually, with 86 Gg-P/year to the continents and 31 Gg-P/year to the oceans and seas. The large value over land is dominated by Africa (52 Gg-P/year) and Asia (24 Gg-P/year), both of which contain large areas of desert and reflect short-range transport from these important source regions. There is a considerable spatial variability over the ocean basins. The North Atlantic, North Pacific, and Indian Oceans dominate the absolute mass of Bio-P deposited to the surface waters accounting for 37%, 19%, and 18% of all Bio-P deposited to the oceans and seas, respectively. In the North Atlantic and Indian Ocean this is correlated with the dust mass deposition (52 and 21%), while the North Pacific accounts for only 11% of global dust supply to surface waters. This apparent enrichment is due to relatively enhanced acid dissolution, which will be discussed in the following section. The global importance of the Mediterranean Sea as a region impacted by dust is clearly demonstrated; it accounts for 5% of the total ocean Bio-P and results in the largest deposited mass per unit area over the oceans ($1.5 \mu\text{g P/m}^2/\text{day}$) and is exceeded on land only by Africa ($4.8 \mu\text{g P/m}^2/\text{day}$) and Asia ($1.5 \mu\text{g P/m}^2/\text{day}$), which are both important dust source regions. The NE Atlantic is also strongly impacted because of its proximity to the Saharan Desert, which exhibits a strong outflow over this region and results in a mean deposition rate of $1.2 \mu\text{g P/m}^2/\text{day}$.

3.2. Acid Dissolution vs Labile P

As observed by Stockdale et al. (2016) mineral dust contains Ap-P and Lab-P that both contribute toward the total fraction of inorganic mineral-borne phosphorus that is potentially bioavailable in marine surface waters. Although measurements reported by Stockdale et al. (2016) find Lab-P present in much smaller quantities than Ap-P, the total mass of Lab-P is directly bioavailable in the surface waters, whereas the Ap-P requires the presence of acid to yield a bioavailable form of P, hereafter referred to as Acid-P. The differences in immediate bioavailability of the two components result in distinct regional differences that can be seen in Figures 3a and 3b. Close to the dust source regions (Saharan, Arabian, Gobi, Patagonian, Kalahari, Great Basin, and Great Australian deserts) the Lab-P dominates the total Bio-P deposition, whereas further away from source regions and over industrialized regions Acid-P dominates; the formation of Acid-P is enhanced as dust accumulates condensed acids during long-range transport and when dust loading is low in




Figure 2. Annual deposition rate of (a) bioavailable phosphorus in $\mu\text{g-P m}^{-2}\text{yr}^{-1}$, (b) dust in $\text{g/m}^2/\text{year}$, and (c) dust-associated H^+ ions in $\text{mol-H}^+ \text{m}^{-2}\text{yr}^{-1}$.

proximity to local pollution sources (thus highlighting areas that are possibly acid-limited in respect to the dissolution of Ap-P). DMS can also play a role in providing a source of acid over the open oceans, especially in the Pacific Ocean, Southern Ocean, and Arctic Ocean (Belviso et al., 2004).

Tables 1 and 2 provide regional information on the annual mean contribution from Lab-P and Acid-P. Of the total 31.2 Gg-P/year of Bio-P deposited to the oceans 16.9 Gg-P/year is from Acid-P and 14.3 Gg P/year from Lab-P. The North Atlantic Ocean, Mediterranean Sea, and the western Indian Ocean, all regions strongly influenced by their proximity to significant dust sources, are dominated by the Lab-P fraction with 94% of the ocean's total Lab-P mass being deposited in these three regions (73, 7, and 14%, respectively). Conversely, the Pacific Ocean, Arctic Ocean, SE Atlantic, South China Sea, and the Southern Ocean, regions further from dust source regions, are dominated by the acid dissolution process; in these regions the contribution of Acid-P to the total Bio-P ranges from 75 to 81%. Although dominated by Lab-P the North Atlantic Ocean, Mediterranean Sea, and western Indian Ocean also receive a considerable mass of Bio-P from Acid-P. Considerable acid dissolution of Ap-P occurs over the Pacific Ocean even though this region receives a very small mass of dust (14% of global oceans dust supply; 25% of global oceans Bio-P supply).

3.3. Spatial Variability of TP Bioavailability

Estimated percentages of TP bioavailability for dust are shown in Figure 3c and Table 2. The simulated TP bioavailability from dust varies from ~10 to 60% and is generally negatively correlated with areas of high dust loading and positively correlated with acid loading. The Lab-P content in dust tends to only vary between 9 and 11% of the TP across the globe (see Table 2). As this pool of phosphorus is deemed directly bioavailable,

Table 1

Absolute Mass of Bioavailable Phosphorus (Total and From Each Source) and Dust Components Deposited to the Surface of Each Region in the Simulated Year

Region	Bioavailable phosphorus deposition			Deposited dust content	
	Total (Bio-P)	From acid dissolution (Acid-P)	From labile pool (Lab-P)	Total phosphorus (TP)	Total dust ^a
Ice sheets					
Greenland	0.03 (18)	0.03 (15)	0.01 (3)	0.06 (30)	0.11 (57)
Antarctic	0.05 (3)	0.04 (3)	0.01 (1)	0.09 (6)	0.12 (8)
Oceans and Seas					
Arctic Ocean	0.5 (36)	0.4 (29)	0.1 (7)	0.9 (66)	1.7 (123)
Pacific Ocean					
NE	2.3 (66)	1.8 (51)	0.5 (15)	5.2 (147)	10.7 (303)
SE	0.8 (14)	0.6 (11)	0.1 (3)	1.5 (26)	3.0 (53)
NW	3.6 (87)	2.7 (65)	0.9 (22)	9.5 (230)	22.1 (535)
SW	1.2 (38)	0.9 (30)	0.2 (8)	2.6 (86)	5.9 (192)
Atlantic Ocean					
NE	6.9 (436)	2.0 (125)	4.9 (310)	48.4 (3041)	103.4 (6497)
SE	2.1 (70)	1.6 (52)	0.5 (18)	5.5 (184)	10.9 (360)
NW	4.7 (181)	2.2 (84)	2.5 (98)	24.5 (943)	50.8 (1959)
SW	1.1 (95)	0.6 (56)	0.4 (39)	4.9 (431)	7.8 (689)
Baltic Sea	0.05 (146)	0.03 (105)	0.01 (40)	0.13 (391)	0.23 (689)
Mediterranean Sea	1.5 (549)	0.5 (177)	1.0 (372)	9.8 (3592)	16.3 (5962)
Indian Ocean					
W	3.5 (89)	1.5 (39)	2.0 (50)	20.2 (516)	42.0 (1074)
E	2.2 (62)	1.4 (39)	0.8 (23)	8.7 (245)	19.7 (559)
South China Sea	0.5 (68)	0.4 (54)	0.1 (14)	1.2 (148)	2.7 (327)
Southern Ocean	0.3 (14)	0.2 (11)	0.1 (3)	0.6 (31)	1.0 (53)
Continents					
Europe	3.5 (364)	1.9 (199)	1.6 (165)	15.4 (1612)	26.9 (2822)
North America	2.0 (87)	1.4 (62)	0.6 (25)	6.1 (270)	10.9 (483)
South America	3.3 (180)	1.4 (79)	1.8 (101)	19.4 (1074)	32.3 (1793)
Africa	51.6 (1738)	3.5 (119)	48.1 (1619)	466.8 (15725)	940.4 (31681)
Asia	23.6 (562)	4.6 (109)	19.0 (453)	198.5 (4725)	425.8 (10138)
Australia	2.3 (282)	0.5 (65)	1.7 (218)	19.9 (2480)	44.5 (5555)
Antarctica	0.05 (3)	0.04 (3)	0.01 (1)	0.09 (6)	0.12 (8)
Totals					
Global total	117 (229)	30 (59)	87 (170)	870 (1695)	1779 (3468)
Ocean total	31.2 (85)	16.9 (46)	14.3 (39)	144 (392)	298 (814)
Land total	86 (595)	13 (92)	73 (503)	726 (5013)	1481 (10225)

Note. All units are in Gg-P/year except for dust which is in Tg/year. Values in parentheses give annual mass per unit area in $\mu\text{g P}/\text{m}^2/\text{year}$ except for dust which is in $\text{mg}/\text{m}^2/\text{year}$.

^aDust in Tg/year ($\text{mg}/\text{m}^2/\text{year}$).

Bio-P in regions of high dust loading will be dominated by the Lab-P component and are likely to be less dependent on changes to acid concentrations. The dominance of Lab-P in high dust loading regions can be seen close to dust source regions over land and in the NE Atlantic Ocean where TP bioavailability (14%) is closely matched with the content of Lab-P in the dust (10%). Conversely, in regions that are not dominated by dust loading, such as the central Pacific and the remote Arctic Ocean, Acid-P has a greater impact and results in a greater TP bioavailability (~50%). Due to the almost constant percentage of Lab-P in dust TP content the variability in TP bioavailability across the globe is controlled by the acid dissolution process; this is apparent from the wide range in percentage contribution of Acid-P in Table 2, which ranges from only 7% over Africa and 29% over the NE Atlantic, to over 77% in the Pacific and 81% over the Arctic Ocean. These findings highlight the importance that acid dissolution has on the deposition of Bio-P from dust to remote regions that have fewer sources of nutrients, such as the Greenland and Antarctic ice sheets, and the Arctic and Southern oceans. These estimates only refer to the bioavailability of dust-borne P. Results from the modeling study of Myriokefalitakis et al. (2016) suggest that dust is the dominant driver of the collective bioavailability of P-containing aerosol species in northern hemisphere oceans, whereas in the southern hemisphere primary biological particles are more pronounced in the South Atlantic and South Pacific Oceans, and sea spray in the Southern Ocean. Therefore, our estimates are likely representative of northern hemisphere oceans, but other sources of P may increasingly drive the observed bioavailability in the high southern hemisphere latitudes.




Figure 3. Annual mean percentage contribution of (a) the acid dissolution process and (b) the labile pool of phosphorus to the total bioavailable phosphorus from dust deposition, and (c) the percentage of the total phosphorus content in dust that is bioavailable upon deposition.

3.4. Sensitivity of Acid Dissolution to Degree of Particle External Mixing

The assumption used in this study is that a given mass of dust particles can be treated as if they are internally mixed, with each dust constituent (Ap-P, Lab-P, and CaCO_3) being present at a constant fraction of each particle's mass; that is, in each grid box every particle will contain the same mass fraction of Ap-P, Lab-P, and CaCO_3 . This is not necessarily important for the Lab-P fraction, but the dissolution of Ap-P into Acid-P occurs simultaneously with the dissolution of CaCO_3 at a rate that is faster than when no CaCO_3 is present.

To understand the importance of the particle mixing assumption, the deposited masses of dust and associated acid were split into distinct components to mimic a population of internally and externally mixed particles of varying concentrations. As previously, all Lab-P is deemed bioavailable, and hence, the deposited Lab-P mass is the same as in the previous experiments. For a given percentage of externally mixed particles the relevant percentage mass of Ap-P is treated separately using equation (2) and assuming no CaCO_3 present, while the remaining mass of Ap-P is assumed internally mixed and treated as per the control experiment using equation (2) with the corresponding mass of CaCO_3 . A third group, consisting of the externally mixed CaCO_3 , is also removed but has no direct impact on the Bio-P and is simply removed along with its associated condensed acid (thereby having an indirect effect on Bio-P). The modeled uptake efficiency of acid onto the mineral particle is independent of the degree of external mixing. The total Bio-P deposited is the sum of Lab-P, internally mixed Acid-P, and externally mixed Acid-P. Externally mixed percentages of 1, 2, 5, 10, 25, 50% and an extreme case of 100% were used to determine the change in total Bio-P deposition to each

Table 2

Annual Means: Percentage of Total-P in Deposited Dust Bioavailable at Surface, Percentage of Bioavailable-P from Acid Dissolution of Apatite, Percentage of Total-P in Dust From Labile-P, and Percentage of Apatite-P in Deposited Dust Dissolved by Acid Dissolution

Region	% of TP in dust bioavailable at surface	% of bio-P from acid dissolution	% of TP in dust from labile pool	% of Ap-P in dust dissolved
Ice sheets				
Greenland	62%	82%	11%	100%
Antarctic	60%	82%	11%	93%
Oceans and Seas				
Arctic Ocean	55%	81%	10%	86%
Pacific Ocean	NE	45%	77%	73%
	SE	52%	81%	84%
	NW	38%	75%	56%
Atlantic Ocean	SW	44%	80%	65%
	NE	14%	29%	9%
	SE	38%	75%	10%
	NW	19%	46%	10%
	SW	22%	59%	9%
Baltic Sea	37%	72%	10%	54%
Mediterranean Sea	15%	32%	10%	10%
Indian Ocean	W	17%	43%	10%
	E	25%	63%	9%
South China Sea	46%	79%	10%	70%
Southern Ocean	45%	79%	10%	74%
Continents				
Europe	23%	55%	10%	25%
North America	32%	71%	9%	45%
South America	17%	44%	9%	16%
Africa	11%	7%	10%	2%
Asia	12%	19%	10%	4%
Australia	11%	23%	9%	5%
Antarctica	60%	82%	11%	93%
Global means				
Global mean	14%	26%	10%	7%
Ocean mean	22%	54%	10%	24%
Land mean	12%	16%	10%	4%

region as a percentage of the control experiment where all particles are assumed to be internally mixed (0% externally mixed). Results are shown in Table 3.

The results show that the Bio-P deposition is very sensitive to the assumptions used for the distribution of components between dust particles. All regions show an increasing deposition rate of Bio-P with an increasing degree of external mixing. An increase from 0 to 5% results in a 9% increase of Bio-P to the oceans, with most of this from regions with large dust loadings where acid dissolution of Ap-P is usually strongly buffered by CaCO_3 . Regions that show less sensitivity are those where acid dissolution is already a dominating process and CaCO_3 is close to exhaustion; in this case the enhanced dissolution rate of Ap-P is already occurring. In reality it is unlikely that CaCO_3 and Ap-P will be exclusively present on different particles, but this provides insight of how sensitive the process is and clearly suggests that this assumption is important for determining the dissolution process of Ap-P and also other minerals that would usually be buffered by the CaCO_3 .

3.5. Evaluation of Model Simulation

Our modeled deposition fluxes, solely from mineral dust, are evaluated against a data set of observed fluxes and in situ aerosol concentrations from ocean cruises and ground-based observations compiled by Vet et al. (2014), which includes data from a number of sources (Baker, French, et al., 2006; Baker, Jickells, et al., 2006; Baker et al., 2010; Carbo et al., 2005; Chen, 2004; Chen & Chen, 2008; Furutani et al., 2010; Herut et al., 1999; Kocak et al., 2005; Koelliker et al., 2004; Markaki et al., 2003; Migon & Sandroni, 1999; Morales-Baquero et al., 2006; NADP/AIRMoN, 2010; Oredalen et al., 2010; Ozsoy, 2003; Rodríguez et al., 2011; Tamatamah et al., 2005; Zhang et al., 2007). Figures 4 and 5a show the observations compared to the modeled deposition fluxes from

Table 3
Sensitivity of Annual Deposited Mass of Bioavailable Phosphorus to Degree of External Mixing of Apatite and Calcium Carbonate (CaCO₃) Content in the Deposited Dust

Region	Percentage of apatite and CaCO ₃ externally mixed in dust						
	1%	2%	5%	10%	25%	50%	100%
Ice sheets							
Greenland	0	0	0	0	0	0	0
Antarctic	0	0	0	1	2	4	5
Oceans and Seas							
Arctic	0	0	1	2	4	7	11
Pacific	NE	0	1	2	4	9	24
	SE	0	0	1	2	4	12
	NW	1	1	4	7	17	33
Atlantic	SW	1	1	3	5	12	24
	NE	3	6	15	31	77	153
	SE	1	1	3	6	15	29
	NW	2	4	10	20	51	102
Baltic	SW	2	3	8	16	39	78
		1	1	3	7	17	33
Mediterranean	3	6	15	29	73	148	221
Indian	W	3	5	13	26	66	132
	E	1	3	7	15	37	74
South China	0	1	2	4	9	19	27
Southern	0	1	2	3	8	16	23
Ocean total	2	4	9	18	44	88	131

Note. All data in % increase compared to control run. Data show percentage increase of bioavailable phosphorus compared to the control run (0% mixing; internally mixed assumption). An externally mixed percentage of 0% implies that apatite and dust are present at the same ratio in all dust particles (therefore internally mixed), and a percentage of 100% implies that apatite and calcium carbonate are exclusively present in different dust particles.

our study. Although limited by observations, the model captures the regional variations in TP: the highest values are found close to source regions, such as the Mediterranean Sea and off the eastern coast of Africa, and lower values in more remote regions, such as the central Pacific. Apparently linear features in Figure 5a for the Pacific coast and Mediterranean occur due to multiple observations being present in the same model grid-box. Standard deviations of model data, shown in Figure 5a, show considerable variability throughout the year. The observational data set contains P from all sources, whereas we only estimate fluxes for mineral dust. The results support the idea that mineral dust is a globally important source of P, but it is worth noting that in regions of low dust loading, such as the Southeast Pacific and Southern Ocean, other sources of P may play a more important role (Myriokefalitakis et al., 2016).

We also compared our Bio-P estimates to field observations in Figures 4 and 5b. The model reproduces the observations within an order of magnitude for much of the Atlantic Ocean and Mediterranean Sea but, as with TP, is underestimating values over the Pacific Ocean and the land; the under-representation over land is likely due to the enhanced role of other sources of P. A key result is that when acid dissolution is removed from the model, shown in Figure 5b using crossed symbols, the Bio-P values are systematically too low compared with measurements. The normalized mean bias (and with respect to the logarithm) without acid dissolution is -0.60 (-0.27) and -0.54 (-0.1) when acid dissolution is included. Our model confirms that acid dissolution in the atmosphere is an important process affecting the deposited flux of Bio-P (through the addition of Acid-P) to the Earth's surface.

Overall TP deposition fluxes to ocean basins and seas are compared to other modeling studies in Table 4. Regional trends show good agreement with other studies: The North Atlantic Ocean is the globally dominant region for deposition, with the North Pacific Ocean and Indian Ocean also important regions for deposition. Our results show good agreement with

Zhang et al. (2015), who similarly only considered mineral dust, in all regions except for the North Atlantic for which the authors estimate an additional 40% deposition of TP. Okin et al. (2011) and Myriokefalitakis et al. (2016) estimate similar fluxes for the Atlantic Ocean but estimate a factor of ~ 2 to 3 greater deposition to the Pacific and Indian oceans. Mahowald et al. (2008) estimate considerably greater fluxes in most regions: the Atlantic Ocean is a factor of 3 greater, and the Pacific and Indian oceans are a factor of ~ 4 greater than our estimates. These differences are largely due to the assumed P content of dust. In our study the P content of the emitted dust was determined using a soil P-content database from Yang et al. (2013), which results in a global mean TP content of 489 ppm per mass of dust. Mahowald et al. (2008) assumed a constant P content of 720 ppm, Okin et al. (2011) use 700 ppm, while Myriokefalitakis et al. (2016) use a soil database (Nickovic et al., 2012) and tune emissions to result in a global mean dust TP content of 880 ppm. Using these alternative global mean contents and our modeled dust fluxes would result in equivalent ocean TP deposition fluxes of 212, 206, and 259 Gg-P/year.

Table 4 also compares Bio-P from this study with other modeling studies. Zhang et al. (2015) provide a good comparison as their study only considers inorganic P from dust and uses a globally constant TP bioavailability for each mineral, experimentally determined in solution with a pH of 2. In contrast, we use a parameterization based on experiments using a full range of conditions, which results in a TP bioavailability that depends on acidity levels and dust loading. In Zhang et al. (2015) the TP bioavailability is $\sim 10\%$ for most of the basins, whereas in this study the bioavailability varies from $\sim 15\%$ in areas of high dust loading to $>45\%$ in regions of high pollution or remote regions only accessible through long-range transport. This results in a global deposition of Bio-P in our study (31.2 Gg-P/year) that is roughly double that of Zhang et al. (2015). Mahowald et al. (2008) estimate Bio-P ocean deposition of 96.5 Gg-P/year, which is roughly 3




Figure 4. Modeled deposition fluxes (contours) and observations (filled circles) from Vet et al. (2014) in $\mu\text{g P/m}^2/\text{year}$ for (a) wet-deposited total P (TP), (b) dry-deposited TP, (c) wet-deposited Bio-P, and (d) dry-deposited Bio-P. Modeled and observational values follow the same color scale.

times our estimate. In their study a constant TP bioavailability of 10% for mineral dust is used. However, the inclusion of other P sources results in a TP bioavailability from all P sources that is greater than 10% in regions where other P sources are important, such as the Pacific and Arctic. The increased TP from other sources, and from the assumed TP content of dust, results in a greater Bio-P deposition estimate than our study.

Myriokefalitakis et al. (2016) estimate 88.7 Gg-P/year Bio-P from inorganic sources only (dust-borne Lab-P, dissolved Ap-P, and volcanic aerosols), which is a factor of ~ 3 greater than our study; this difference arises mainly through differences in the TP content of dust, as well as a different modeled acid dissolution process, and through contributions from volcanic aerosols (6 Gg-P/year predominantly to the Pacific Ocean). If we apply the higher dust TP and Lab-P content from Myriokefalitakis et al. (2016), it increases our global Bio-P estimate by 16 Gg-P/year. The final difference occurs through the treatment of the Acid-P production; we estimate that 30 Gg-P/year is deposited globally as Acid-P, compared to 144 Gg-P/year by Myriokefalitakis et al. (2016). Part of this difference in Acid-P is due to the assumption used in Myriokefalitakis et al. (2016) that the TP content of dust is composed of Lab-P and Ap-P only, whereas other constituents are included in our soil database (e.g., Lab-P and Ap-P make up $\sim 50\%$ of TP in the Sahara Desert). Assuming our deposited dust is only composed of Lab-P and Ap-P would provide a factor of 3 increase in Ap-P availability, which as shown by our sensitivity study (Table S2) could have a considerable impact on Bio-P deposition. The final differences in Acid-P production occur in the different representations of acid dissolution: in our study the empirically parameterized process is dependent on the absolute number of H^+ ions from H_2SO_4 and HNO_3 uptake and calculated offline after deposition, whereas in Myriokefalitakis et al. (2016) an online aerosol thermodynamic equilibrium framework is used to determine conditions for use in a kinetic model. The added sources of atmospheric acid and detailed treatment of the particle's aqueous composition result in greater acid dissolution than the simple experimentally based method used in our study. However, despite these differences it is an interesting result that the two methods produce similar spatial patterns but of different magnitudes. Assessing these differences and identifying the more appropriate method are beyond the scope of this study but merit further attention.




Figure 5. Data as in Figure 4 but shown as scatter plots with observations from Vet et al. (2014) on the x axis and modeled fluxes from this study on the y axis. (a) Data for total P (TP) fluxes and (b) for Bio-P fluxes. Different colored circles depict different regions. Vertical lines in (a) show the standard deviation of modeled deposition flux at that location from the simulated year (not repeated in (b) for clarity). Crosses in (b) show the modeled deposition flux of Bio-P without simulated acid dissolution of apatite.

4. Discussion

There is large spatial variability in the amount of Bio-P supplied to the surface ocean. The largest flux of dust and of Bio-P, predominantly in the form of Lab-P, is supplied to the NE Atlantic Ocean; this area is known to have a high dust flux from the world's major source of mineral dust, the Sahara Desert. The modeling results suggest that this area not only has the highest total input of Bio-P (6.9 Gg-P/year) but also a high input of Bio-P per unit area (436 $\mu\text{g P/m}^2/\text{year}$; second to the Mediterranean Sea with 549 $\mu\text{g P/m}^2/\text{year}$). Neuer et al. (2004) suggest that the pulses of Saharan dust seem to exert a feast or famine effect on phytoplankton export from this region. A pulsing that may be linked to the observed phytoplankton export is apparent in the modeled fluxes (Figures S3 to S6), which vary between 0.2 and 0.6 Gg-P per month in the NE Atlantic and between 0.4 and 0.9 Gg-P per month in the NW Atlantic, another region with considerable deposition of Bio-P. In these regions, Saharan dust plumes transported over the North Atlantic meet North American air masses, which are enriched in anthropogenic combustion products. Sedwick et al. (2007) identified an increase in soluble Fe delivered to the offshore NW Atlantic caused by the North American polluted air masses. Here we show based on our modeling that these air masses also increase the amount of Acid-P from the mineral dust transported across the Atlantic from the Sahara, especially during the summer months when dust transport and acid pollution are both relatively enhanced.

The total Bio-P supplied to the ocean is the sum of contributions from Lab-P and Acid-P. The location with the highest mass of deposited Bio-P from Acid-P is the NW Pacific (3.6 Gg-P/year). Our data (Table 1) are averaged over the whole NW Pacific area; however, the impact is greatest in the area directly under the plume from the Gobi and Taklamakan deserts, which passes over the highly polluted air in East Asia (Jaffe et al., 1999) and out into the NW Pacific (Chen et al., 2017). This plume is rather narrowly confined to $\sim 40^\circ\text{N}$ but has a large effect within that plume as shown in Figures S3 and S4. We see the strongest peak during the Northern Hemisphere spring (March–April–May; Figure S6), which is consistent with the findings of Chen et al. (2017). Maki et al. (2016) find that Asian dust is a major nutrient source to the area of the NW Pacific under the plume though their study concentrates on the flux of atmospheric nitrate to this N limited region. As argued below the greatly increased flux of Bio-P from the dust plume will reduce the potential effect of switching to P limitation

and may also increase the N_2 fixation in the surface waters (Jickells & Moore, 2015) as well as increasing the total primary productivity over longer timescales (Tyrrell, 1999).

The location with the highest annual mass of Bio-P per unit area ($\sim 550 \mu\text{g P/m}^2/\text{year}$) is the Mediterranean Sea. This area is particularly sensitive to the addition of extra Bio-P since both the western and particularly the eastern Mediterranean are P limited (Krom et al., 1991; Moutin & Raimbault, 2002; Powley et al., 2017). Any addition of P available to the phytoplankton will rapidly be taken up into the P-starved water (Krom et al., 2005) and be translated into extra primary productivity. Microcosm experiments have shown the importance of dust inputs to increasing primary productivity particularly during the summer when there is minimal nutrient supply from below (Laghdass et al., 2012; Ridame et al., 2014). Krom et al. (2010) estimated that Lab-P from atmospheric sources provided 30% of the nonmarine external P supplied to the eastern Mediterranean, while Powley et al. (2017), using a mass balance model, estimated that 49% of the nonmarine external supply to the entire Mediterranean was from atmospheric supply. African dust outbreaks are estimated by Pey et al. (2013) to occur over the Mediterranean over 20% of annual days. These plumes of mineral dust, along with locally derived sources, meet air polluted with acidic gases from Europe including air that has passed over

Table 4
Modeled Deposition Fluxes of TP and Bio-P to Different Ocean Basins in Gg-P/year

	Mahowald et al. (2008)		Okin et al. (2011)		Zhang et al. (2015) ^a		Myriokefalitakis et al. (2016)			This study ^a	
	TP	Bio-P	TP	Bio-P	TP	Bio-P sol-1	TP	Bio-P DP	Bio-P DIP ^b	TP	Bio-P
Atlantic	231	35.8 (15)	128	–	112	9.6 (9)	110	54.9 (50)	36.6	83	14.8 (18)
N Atlantic	190	27.9 (15)	–	–	103	8.8 (9)	96	44.2 (46)	32.3	73	11.6 (16)
S Atlantic	42	7.9 (19)	–	–	9	0.8 (9)	14	10.6 (74)	4.4	10	3.2 (30)
Pacific	85	28 (33)	68	–	18	1.8 (9)	59	48.4 (81)	26.6	20	8.4 (42)
N Pacific	63	22 (35)	–	–	18	1.7 (10)	47	38.0 (80)	23.9	15	5.9 (40)
S Pacific	22	6 (27)	–	–	1	0.07 (8)	12	10.5 (86)	2.7	5	2.5 (47)
Indian	149	20.3 (14)	73	–	37	3.8 (10)	56	37.3 (66)	20.7	29	5.7 (20)
Southern	12	1.7 (14)	32	–	0.2	0.01 (7)	2	1.5 (82)	0.2	1	0.5 (46)
Arctic	7	2.2 (32)	–	–	1	0.1 (10)	4	3.3 (74)	1.7	1	0.5 (55)
Mediterranean	54	5.7 (11)	–	–	11	1.1 (10)	7	4.4 (59)	2.7	10	1.5 (15)
Antarctic ice	–	–	–	–	0.1	0.01 (9)	0.1	0.0 (23)	0.01	0.1	0.1 (60)
Greenland ice	–	–	–	–	5	0.5 (9)	0.3	0.2 (62)	0.1	0.1	0.03 (62)
Ocean total	558	96.5 (17)	320	–	180	16.3 (9)	240	150.2 (63)	88.7	144	31.2 (22)
Land total	832	143.5 (17)	–	–	–	–	1051	303 (29)	160	726	86 (12)

Note. Figures in parentheses correspond to the TP bioavailability (%) upon deposition. Bio-P data from Zhang et al. (2015) are presented for their “sol-1” data set (please refer to the study for details). Values for both total dissolved phosphorus (DP) and dissolved inorganic phosphorus (DIP) from Myriokefalitakis et al. (2016) are included. Values from Zhang et al. (2015) and this study are only for phosphorus from inorganic mineral dust.

^aMineral dust only. ^bDissolved inorganic P from dust and volcanic emissions only.

cities such as Athens (Nenes et al., 2011); this results in extra Acid-P, and thus Bio-P, being produced. The modeling results are compatible with the predictions made by Nenes et al. (2011) concerning the effect of acid processing on mineral aerosol over the Mediterranean.

There is also a considerable contribution from Acid-P in both the western and eastern Indian Ocean. This is an area where dust from local desert regions such as the Arabian Peninsula and the Thar desert in India meet with air masses polluted from highly populated areas in South and South-East Asia (Lelieveld et al., 2001) and pollution from biomass burning over Central and South-eastern Africa and South Asia (Sinha et al., 2004). It has been suggested that atmospheric aerosols including dust particles, as well as biomass burning and other anthropogenic sources, are increasing the flux of dissolved nutrients (including phosphate) to the Bay of Bengal (Srinivas & Sarin, 2015). Rengarajan and Sarin (2004) suggested that aerosols that have been involved in interactions between mineral dust and pollution also impacted the Arabian Sea; they identified its effect on Fe and NO_x. Here we suggest that it also affects the Acid-P.

It is generally assumed that any potential Bio-P (i.e., mineral apatite and/or Fe-bound P) as well as Lab-P that lands on most parts of the terrestrial system will be consumed by plants. Plants and their accompanying mycorrhizal fungi are evolved to extract this key nutrient from such minerals present in the soil (Smits et al., 2012). The only exception to this might be areas of snow, such as Greenland and the Arctic, which host algal communities, which are important for climate change due to their impact on surface albedo (Lutz et al., 2016). It has been reported that microbes in arctic glaciers live in P-deficient environments (Stibal et al., 2009), which suggests that atmospheric input of Bio-P to these key regions may play an important role on local algal and microbial activity. We estimate that the amount of Bio-P delivered to the major ice sheets of the world (Greenland and Antarctica) is 0.08 Gg-P/year of which the majority (~80%) is produced by acid dissolution.

Modeling studies routinely use a constant value for TP bioavailability, which typically ranges from ~10 to ~15%. From the soil P database used in our study we show that TP bioavailability is at least 10% due to Lab-P alone and is considerably greater once the Acid-P contribution is included. For inorganic Bio-P from mineral dust we estimate global mean TP bioavailability of 22 and 12% for oceans and land, respectively. For the ocean basins, we estimate the following: Pacific Ocean (42%), Atlantic Ocean (18%), Indian Ocean (20%), Mediterranean (15%), Southern Ocean (46%), and Arctic Ocean (55%). Other studies such as Myriokefalitakis et al. (2016) show similar increases in TP bioavailability caused by Acid-P. These calculations suggest that while the spatial variability of TP and Bio-P are similar (Table 4), the absolute fluxes vary considerably to a large extent because of different values for TP in the source material. Uncertainties in current soil databases primarily arise from a lack of observations and sampling of under-represented soil types, including

species that contain phosphorus (Nickovic et al., 2012; Yang et al., 2013); the result is poor spatial coverage and a lack of process-level understanding that yields considerable uncertainties. The sensitivity experiments (Tables 3 and S2) highlight the clear need for improved estimates of dust-borne P components at source.

Krishnamurthy et al. (2010) calculated that the addition of atmospheric P to the global ocean will increase the total carbon uptake by 0.12% based on an estimated 8.3 Gg-P/year Bio-P deposited to the oceans. However, the authors assumed a globally constant dust TP content of 1050 ppm and a globally constant TP bioavailability of 15%. In comparison, our model estimates Bio-P deposition from inorganic mineral dust to be ~400% of this value (31.2 Gg-P/year) with TP bioavailability ranging from ~15 to ~50%. Our enhanced Bio-P deposition estimate suggests that the total carbon uptake could be even greater, if the apatite and CaCO_3 composition in the mineral dust population were externally mixed. Assuming that all Bio-P is taken up by phytoplankton, and applying the Redfield ratio, our results suggest that the atmospheric input of Bio-P from mineral dust may account for an uptake of 1.3 Tg-C/year, with 0.7 Tg-C/year from acid-dissolution of mineral dust. If the mineral dust Ap-P/ CaCO_3 content is deemed 100% externally mixed, then the total uptake would be 3.0 Tg-C/year.

In non-Fe-limited areas where the system is N limited in the short term, it has been shown that most atmospheric input has a high bioavailable N:P ratio and an excess of Fe. It has been suggested that the input of Fe and N causes the systems to switch toward P limitation (Jickells & Moore, 2015). The increased Bio-P shown to be formed by these atmospheric acid processes linked mainly to anthropogenic pollution will slow this process down particularly in areas like the NW Atlantic (Chien et al., 2016). This study suggests that the abundant acid gases produced during volcanic eruptions are likely to have a short term but possibly dramatic effect on the supply of Bio-P to the ocean. Previous work has been confined to the actual Bio-P produced within the volcanic plume itself, which is rather small (e.g., Mahowald et al., 2008). Here we predict that a more important effect might be to increase the flux of Bio-P to the ocean due to the interaction of acid gases from the volcano with existing mineral particles in the atmosphere.

Stoichiometric proxies have been used to estimate nitrogen fixation rates in oceanic surface waters. Deutsch et al. (2007) calculated a parameter ($P^* = \text{PO}_4^{3-} - \text{NO}_3^-/16$) and suggested that regions with high P^* are also regions of the ocean with high N_2 fixation rates and vice versa. It is noticeable that the three regions of the global ocean with low values of P^* in their calculations (north central Atlantic, north western Pacific, and Mediterranean) are also the areas with the predicted highest fluxes of Acid-P. Superficially, the addition of extra bio-P should increase P^* ; however, a large fraction of the acid, which produces acid-P, is nitric acid. This is apparent in the eastern Mediterranean where the atmospheric N:P ratio has been observed as high as 105:1 (Markaki et al., 2010). This is an area of the ocean with very low N_2 fixation rates (Yogev et al., 2011). By contrast, the N_2 fixation rate of the north central Atlantic is relatively high because the high flux of Saharan dust contains abundant Fe and P, which are required by N_2 fixing organisms (Mills et al., 2004). It is unclear what overall impact the external supply of bio-P from mineral dust has on N_2 fixation rates and P^* ; however, our results provide global estimates that may help to improve our understanding in the future.

5. Summary

In this study a parameterization for acid-dissolution of mineral apatite, developed using results presented by Stockdale et al. (2016), was incorporated into a global aerosol model (GLOMAP; Mann et al., 2010) with a global soil P database (Yang et al., 2013) to model the atmospheric flux of inorganic bioavailable P from mineral dust. We estimate that 870 Gg-P/year of inorganic TP associated with dust is deposited globally, with 726 Gg-P/year to the land and 144 Gg-P/year to the oceans. Our model is able to discriminate between the leachable (labile) pool of phosphorus, which is present upon emission and the dissolved apatite pool, which is a result of the simultaneous acid-dissolution of apatite and calcium-carbonate; the two pools combined represent the atmospheric flux of bioavailable phosphorus (Bio-P) from dust.

We estimate a global flux of 31 Gg-P/year Bio-P to the oceans with 14.3 Gg-P/year from the labile pool (Lab-P) and 16.9 Gg-P/year from the acid-dissolved pool (Acid-P). The acid dissolution of mineral dust increases supply of Bio-P to the oceans by 120%, showing the importance of the acid dissolution process on global Bio-P fluxes. We identify the Mediterranean Sea, North Atlantic Ocean, and North Pacific Ocean as particularly important regions for dust-borne Bio-P deposition.

Our modeling results show that the percentage of deposited dust TP that is in a bioavailable form ranges from ~10 to ~50%; however, generally, it has been assumed in modeling studies that the percentage of mineral dust TP bioavailability is globally constant, with values between 10 and 15% commonly used. We use a similar approach to Myriokefalitakis et al. (2016) to show that although the labile fraction of TP is globally constant at ~10%, the acid-dissolved pool increased the mean TP bioavailability over oceans to 22% with considerable variation between ocean basins: Pacific Ocean (42%), Atlantic Ocean (18%), Indian Ocean (20%), and Mediterranean (15%). This variability again highlights the impact that the acid-dissolution process has on the global flux of Bio-P from dust.

The world's largest dust sources all have relatively enriched levels of dust-P pools but show variability between sources. We therefore advise against using global constant values for dust-P pools and recommend using global databases of soil-P content. In our study we used a pedogenesis-based soil-P database from Yang et al. (2013), which resulted in a global mean deposited dust TP content of 489 ppm with 49 ppm from Lab-P and 243 ppm from Ap-P, with all pools displaying roughly 1 order of magnitude variability across the globe. Comparing these results with other modeling studies and results from a series of sensitivity tests highlighted the affect that the assumed dust mineralogical content has on the Bio-P flux. It is noted that there is no consensus on the global mean deposited dust TP and previous studies have used higher values (e.g., 880 ppm by Myriokefalitakis et al., 2016, and 720 ppm by Mahowald et al., 2008).

From our sensitivity study we have identified that treating the dust population as an externally rather than internally mixed population (i.e., assuming that the population exhibits particle-to-particle diversity in apatite and CaCO₃ content) results in a considerable increase in global Bio-P flux (44% increase for a 25% externally mixed population). It is currently not known which treatment is appropriate and requires more research.

Our results confirm the importance of acid processes in the atmosphere in increasing the flux of Bio-P to the global ocean as suggested by Nenes et al. (2011). The effect is spatially and temporally variable, and it is suggested that increased Bio-P can result in regionally important changes in biogeochemical processes such as nutrient limitation, nitrogen fixation rates, and carbon uptake.

Acknowledgments

This research was funded by the Leverhulme Trust Research Project grant RPG 406, entitled "Understanding the Delivery of phosphorus nutrient to the oceans." K. S. C. acknowledges funding from the EU CRESCENDO project under grant agreement 641816. The authors wish to thank T. Nenes, M. Kanakidou, and S. Myriokefalitakis for useful discussions and for providing data used in the comparison to Myriokefalitakis et al. (2016). The authors also wish to thank R. Vet for providing the observational data set used; this source is listed in the references. Other data used are listed in the references, tables, and supporting information.

References

- Atkinson, J. D., Murray, B. J., Woodhouse, M. T., Whale, T. F., Baustian, K. J., Carslaw, K. S., et al. (2013). The importance of feldspar for ice nucleation by mineral dust in mixed-phase clouds. *Nature*, *498*(7454), 355–358. <https://doi.org/10.1038/nature12278>
- Baker, A. R., French, M., & Linge, K. L. (2006). Trends in aerosol nutrient solubility along a west–east transect of the Saharan dust plume. *Geophysical Research Letters*, *33*, L07805. <https://doi.org/10.1029/2005GL024764>
- Baker, A. R., Jickells, T. D., Witt, M., & Linge, K. L. (2006). Trends in the solubility of iron, aluminium, manganese and phosphorus in aerosol collected over the Atlantic Ocean. *Marine Chemistry*, *98*(1), 43–58. <https://doi.org/10.1016/j.marchem.2005.06.004>
- Baker, A. R., Lesworth, T., Adams, C., Jickells, T. D., & Ganzeveld, L. (2010). Estimation of atmospheric nutrient inputs to the Atlantic Ocean from 50°N to 50°S based on large-scale field sampling: Fixed nitrogen and dry deposition of phosphorus. *Global Biogeochemical Cycles*, *24*, GB3006. <https://doi.org/10.1029/2009GB003634>
- Belviso, S., Moulin, C., Bopp, L., & Stefels, J. (2004). Assessment of a global climatology of oceanic dimethylsulfide (DMS) concentrations based on SeaWiFS imagery (1998–2001). *Canadian Journal of Fisheries and Aquatic Sciences*, *61*(5), 804–816. <https://doi.org/10.1139/f04-001>
- Brahney, J., Mahowald, N., Ward, D. S., Ballantyne, A. P., & Neff, J. C. (2015). Is atmospheric phosphorus pollution altering global alpine Lake stoichiometry? *Global Biogeochemical Cycles*, *29*, 1369–1383. <https://doi.org/10.1002/2015gb005137>
- Browse, J., Carslaw, K. S., Arnold, S. R., Pringle, K., & Boucher, O. (2012). The scavenging processes controlling the seasonal cycle in Arctic sulfate and black carbon aerosol. *Atmospheric Chemistry and Physics*, *12*(15), 6775–6798. <https://doi.org/10.5194/acp-12-6775-2012>
- Carbo, P., Krom, M. D., Homoky, W. B., Benning, L. G., & Herut, B. (2005). Impact of atmospheric deposition on N and P geochemistry in the southeastern Levantine basin. *Deep Sea Research Part II: Topical Studies in Oceanography*, *52*(22–23), 3041–3053. <https://doi.org/10.1016/j.dsr2.2005.08.014>
- Chen, H.-Y., & Chen, L.-D. (2008). Importance of anthropogenic inputs and continental-derived dust for the distribution and flux of water-soluble nitrogen and phosphorus species in aerosol within the atmosphere over the East China Sea. *Journal of Geophysical Research*, *113*, D11303. <https://doi.org/10.1029/2007JD009491>
- Chen, S., Huang, J., Kang, L., Wang, H., Ma, X., He, Y., et al. (2017). Emission, transport, and radiative effects of mineral dust from the Taklimakan and Gobi deserts: Comparison of measurements and model results. *Atmospheric Chemistry and Physics*, *17*(3), 2401–2421. <https://doi.org/10.5194/acp-17-2401-2017>
- Chen, Y. (2004). Sources and Fate of Atmospheric Nutrients over the Remote Oceans and Their Role on Controlling Marine Diazotrophic Microorganisms, (PhD dissertation). University of Maryland, College Park.
- Chien, C.-T., Mackey, K. R. M., Dutkiewicz, S., Mahowald, N. M., Prospero, J. M., & Paytan, A. (2016). Effects of African dust deposition on phytoplankton in the western tropical Atlantic Ocean off Barbados. *Global Biogeochemical Cycles*, *30*, 716–734. <https://doi.org/10.1002/2015gb005334>
- Chipperfield, M. P. (2006). New version of the TOMCAT/SLIMCAT off-line chemical transport model: Intercomparison of stratospheric tracer experiments. *Quarterly Journal of the Royal Meteorological Society*, *132*(617), 1179–1203. <https://doi.org/10.1256/qj.05.51>

- Dentener, F., Kinne, S., Bond, T., Boucher, O., Cofala, J., Generoso, S., et al. (2006). Emissions of primary aerosol and precursor gases in the years 2000 and 1750 prescribed data-sets for AeroCom. *Atmospheric Chemistry and Physics*, 6(12), 4321–4344. <https://doi.org/10.5194/acp-6-4321-2006>
- Deutsch, C., Sarmiento, J. L., Sigman, D. M., Gruber, N., & Dunne, J. P. (2007). Spatial coupling of nitrogen inputs and losses in the ocean. *Nature*, 445(7124), 163–167. <https://doi.org/10.1038/nature05392>
- Diehl, T., Heil, A., Chin, M., Pan, X., Streets, D., Schultz, M., & Kinne, S. (2012). Anthropogenic, biomass burning, and volcanic emissions of black carbon, organic carbon, and SO₂ from 1980 to 2010 for hindcast model experiments. *Atmospheric Chemistry and Physics Discussions*, 12(9), 24,895–24,954. <https://doi.org/10.5194/acpd-12-24895-2012>
- Eijsink, L. M., Krom, M. D., & Herut, B. (2000). Speciation and burial flux of phosphorus in the surface sediments of the eastern Mediterranean. *American Journal of Science*, 300(6), 483–503. <https://doi.org/10.2475/ajs.300.6.483>
- Fuchs, N. A., & Sutugin, A. G. (1971). High-dispersed aerosols. In J. R. Brock (Ed.), *Topics in current aerosol research* (p. 1–60). Oxford: Pergamon.
- Furutani, H., Meguro, A., Iguchi, H., & Uematsu, M. (2010). Geographical distribution and sources of phosphorus in atmospheric aerosol over the North Pacific Ocean. *Geophysical Research Letters*, 37, L03805. <https://doi.org/10.1029/2009GL041367>
- Graham, W. F., & Duce, R. A. (1982). The atmospheric transport of phosphorus to the western North Atlantic. *Atmospheric Environment*, 16(5), 1089–1097. [https://doi.org/10.1016/0004-6981\(82\)90198-6](https://doi.org/10.1016/0004-6981(82)90198-6)
- Herut, B., Krom, M. D., Pan, G., & Mortimer, R. (1999). Atmospheric input of nitrogen and phosphorus to the Southeast Mediterranean: Sources, fluxes, and possible impact. *Limnology and Oceanography*, 44(7), 1683–1692. <https://doi.org/10.4319/lo.1999.44.7.1683>
- Herut, B., Zohary, T., Krom, M. D., Mantoura, R. F. C., Pitta, P., Psarra, S., et al. (2005). Response of East Mediterranean surface water to Saharan dust: On-board microcosm experiment and field observations. *Deep Sea Research Part II: Topical Studies in Oceanography*, 52(22–23), 3024–3040. <https://doi.org/10.1016/j.dsr2.2005.09.003>
- Huneeeus, N., Schulz, M., Balkanski, Y., Griesfeller, J., Prospero, J., Kinne, S., et al. (2011). Global dust model intercomparison in AeroCom phase I. *Atmospheric Chemistry and Physics*, 11(15), 7781–7816. <https://doi.org/10.5194/acp-11-7781-2011>
- Jaffe, D., Anderson, T., Covert, D., Kotchenruther, R., Trost, B., Danielson, J., et al. (1999). Transport of Asian air pollution to North America. *Geophysical Research Letters*, 26, 711–714. <https://doi.org/10.1029/1999GL900100>
- Jickells, T., & Moore, C. M. (2015). The importance of atmospheric deposition for ocean productivity. *Annual Review of Ecology, Evolution, and Systematics*, 46(1), 481–501. <https://doi.org/10.1146/annurev-ecolsys-112414-054118>
- Kettle, A. J., Andreae, M. O., Amouroux, D., Andreae, T. W., Bates, T. S., Berresheim, H., et al. (1999). A global database of sea surface dimethylsulfide (DMS) measurements and a procedure to predict sea surface DMS as a function of latitude, longitude, and month. *Global Biogeochemical Cycles*, 13, 399–444. <https://doi.org/10.1029/1999GB900004>
- Kocak, M., Kubilay, N., Herut, B., & Nimmo, K. (2005). Dry atmospheric fluxes of trace metals (Al, Fe, Mn, Pb, Cd, Zn, Cu) over the Levantine Basin: A refined assessment. *Atmospheric Environment*, 39(38), 7330–7341. <https://doi.org/10.1016/j.atmosenv.2005.09.010>
- Koelliker, Y., Totten, L. A., Gigliotti, C. L., Offenberg, J. H., Reinfelder, J. R., Zhuang, Y., & Eisenreich, S. J. (2004). Atmospheric wet deposition of total phosphorus in New Jersey. *Water, Air, & Soil Pollution*, 154(1–4), 139–150. <https://doi.org/10.1023/B:WATE.0000022952.12577.c5>
- Krishnamurthy, A., Moore, J. K., Mahowald, N., Luo, C., & Zender, C. S. (2010). Impacts of atmospheric nutrient inputs on marine biogeochemistry. *Journal of Geophysical Research*, 115, G01006. <https://doi.org/10.1029/2009JG001115>
- Krom, M. D., Emeis, K. C., & Van Cappellen, P. (2010). Why is the eastern Mediterranean phosphorus limited? *Progress in Oceanography*, 85(3–4), 236–244. <https://doi.org/10.1016/j.poccean.2010.03.003>
- Krom, M. D., Kress, N., Brenner, S., & Gordon, L. I. (1991). Phosphorus limitation of primary productivity in the eastern Mediterranean Sea. *Limnology and Oceanography*, 36(3), 424–432. <https://doi.org/10.4319/lo.1991.36.3.0424>
- Krom, M. D., Thingstad, T. F., Brenner, S., Carbo, P., Drakopoulos, P., Fileman, T. W., et al. (2005). Summary and overview of the CYCLOPS P addition Lagrangian experiment in the eastern Mediterranean. *Deep Sea Research Part II: Topical Studies in Oceanography*, 52(22–23), 3090–3108. <https://doi.org/10.1016/j.dsr2.2005.08.018>
- Laghdass, M., Catala, P., Caparros, J., Oriol, L., Lebaron, P., & Obermosterer, I. (2012). High contribution of SAR11 to microbial activity in the north West Mediterranean Sea. *Microbial Ecology*, 63(2), 324–333. <https://doi.org/10.1007/s00248-011-9915-7>
- Lelieveld, J., Crutzen, P. J., Ramanathan, V., Andreae, M. O., Brenninkmeijer, C. A. M., Campos, T., et al. (2001). The Indian Ocean Experiment: Widespread air pollution from South and Southeast Asia. *Science*, 291(5506), 1031–1036. <https://doi.org/10.1126/science.1057103>
- Lutz, S., Anesio, A. M., Raiswell, R., Edwards, A., Newton, R. J., Gill, F., & Benning, L. G. (2016). The biogeography of red snow microbiomes and their role in melting arctic glaciers. *Nature Communications*, 7, 11968. <https://doi.org/10.1038/ncomms11968>
- Mahowald, N., Jickells, T. D., Baker, A. R., Artaxo, P., Benitez-Nelson, C. R., Bergametti, G., et al. (2008). Global distribution of atmospheric phosphorus sources, concentrations and deposition rates, and anthropogenic impacts. *Global Biogeochemical Cycles*, 22, GB4026. <https://doi.org/10.1029/2008GB003240>
- Maki, T., Ishikawa, A., Mastunaga, T., Pointing, S. B., Saito, Y., Kasai, T., et al. (2016). Atmospheric aerosol deposition influences marine microbial communities in oligotrophic surface waters of the western Pacific Ocean. *Deep-Sea Research*, 118, 37–45. <https://doi.org/10.1016/j.dsr.2016.10.002>
- Mann, G. W., Carslaw, K. S., Spracklen, D. V., Ridley, D. A., Manktelow, P. T., Chipperfield, M. P., et al. (2010). Description and evaluation of GLOMAP-mode: A modal global aerosol microphysics model for the UKCA composition-climate model. *Geoscientific Model Development*, 3(2), 519–551. <https://doi.org/10.5194/gmd-3-519-2010>
- Markaki, Z., Lojze-Pilot, M. D., Violaki, K., Benyahya, L., & Mihalopoulos, N. (2010). Variability of atmospheric deposition of dissolved nitrogen and phosphorus in the Mediterranean and possible link to the anomalous seawater N/P ratio. *Marine Chemistry*, 120(1–4), 187–194. <https://doi.org/10.1016/j.marchem.2008.10.005>
- Markaki, Z., Oikonomou, K., Kocak, M., Kouvarakis, G., Chaniotaki, A., Kubilay, N., & Mihalopoulos, N. (2003). Atmospheric deposition of inorganic phosphorus in the Levantine Basin, eastern Mediterranean: Spatial and temporal variability and its role in seawater productivity. *Limnology and Oceanography*, 48(4), 1557–1568. <https://doi.org/10.4319/lo.2003.48.4.1557>
- Migon, C., & Sandroni, V. (1999). Phosphorus in rainwater: Partitioning inputs and impact on the surface coastal ocean. *Limnology and Oceanography*, 44(4), 1160–1165. <https://doi.org/10.4319/lo.1999.44.4.1160>
- Mills, M. M., Ridame, C., Davey, M., La Roche, J., & Geider, R. J. (2004). Iron and phosphorus co-limit nitrogen fixation in the eastern tropical North Atlantic. *Nature*, 429(6989), 292–294. <https://doi.org/10.1038/nature02550>
- Morales-Baquero, R., Pulido-Villena, E., & Reche, I. (2006). Atmospheric inputs of phosphorus and nitrogen to the southwest Mediterranean region: Biogeochemical responses of high mountain lakes. *Limnology and Oceanography*, 51(2), 830–837. <https://doi.org/10.4319/lo.2006.51.2.0830>
- Moutin, T., & Raimbault, P. (2002). Primary production, carbon export and nutrients availability in western and eastern Mediterranean Sea in early summer 1996 (MINOS cruise). *Journal of Marine Systems*, 33–34, 273–288. [https://doi.org/10.1016/S0924-7963\(02\)00062-3](https://doi.org/10.1016/S0924-7963(02)00062-3)

- Myrriokefalitakis, S., Nenes, A., & Baker, A. R. (2016). Mihalopoulos, N., and Kanakidou, M.: Bioavailable atmospheric phosphorus supply to the global ocean: A 3-D global modeling study. *Biogeosciences*, 13(24), 6519–6543. <https://doi.org/10.5194/bg-13-6519-2016>
- NADP/AIRMoN: (Atmospheric Integrated Research Monitoring Network) (2010). National Atmospheric Deposition Program (NADP) Office, Illinois State Water Survey, 2204 Griffith Dr., Champaign, IL. Retrieved from <http://nadp.sws.uiuc.edu/AIRMoN/>
- Nenes, A., Krom, M. D., Mihalopoulos, N., Van Cappellen, P., Shi, Z., Bougiatioti, A., et al. (2011). Atmospheric acidification of mineral aerosols: A source of bioavailable phosphorus for the oceans. *Atmospheric Chemistry and Physics*, 11(13), 6265–6272. <https://doi.org/10.5194/acp-11-6265-2011>
- Neuer, S., Torres-Padrón, M. E., Gelado-Caballero, M. D., Rueda, M. J., Hernández-Brito, J., Davenport, R., & Wefer, G. (2004). Dust deposition pulses to the eastern subtropical North Atlantic gyre: Does ocean's biogeochemistry respond? *Global Biogeochemical Cycles*, 18, GB4020. <https://doi.org/10.1029/2004GB002228>
- Nickovic, S., Vukovic, A., Vujadinovic, M., Djurdjevic, V., & Pejanovic, G. (2012). Technical note: High-resolution mineralogical database of dust-productive soils for atmospheric dust modeling. *Atmospheric Chemistry and Physics*, 12(2), 845–855. <https://doi.org/10.5194/acp-12-845-2012>
- Nightingale, P. D., Malin, G., Law, C. S., Watson, A. J., Liss, P. S., Liddicoat, M. I., et al. (2000). In situ evaluation of air-sea gas exchange parameterizations using novel conservative and volatile tracers. *Global Biogeochemical Cycles*, 14, 373–387. <https://doi.org/10.1029/1999GB900091>
- Okin, G. S., Baker, A. R., Tegen, I., Mahowald, N. M., Dentener, F. J., Duce, R. A., et al. (2011). Impacts of atmospheric nutrient deposition on marine productivity: Roles of nitrogen, phosphorus, and iron. *Global Biogeochemical Cycles*, 25, GB2022. <https://doi.org/10.1029/2010GB003858>
- Oredalen, T., Aas, W., and Maenhaut, W. (2010). Atmospheric dry and wet deposition of phosphorus in southern Norway. In: *Data obtained from the World Meteorological Organization Scientific Advisory Group of Precipitation Chemistry Workshop*. Berg-en-Dal, South Africa, 15–20 March 2010.
- Ozsoy, T. (2003). Atmospheric wet deposition of soluble macro-nutrients in the Cilician Basin, north-eastern Mediterranean Sea. *Journal of Environmental Monitoring*, 5(6), 971–976. <https://doi.org/10.1039/B309636J>
- Parkhurst, D. L., & Appelo, C. A. J. (2013). Description of input and examples for PHREEQC version 3—A computer program for speciation, batch-reaction, one-dimensional transport, and inverse geochemical calculations. U.S. Geological Survey Techniques and Methods, Book 6.
- Pey, J., Querol, X., Alastuey, A., Forastiere, F., & Stafoggia, M. (2013). African dust outbreaks over the Mediterranean Basin during 2001–2011: PM₁₀ concentrations, phenomenology and trends, and its relation with synoptic and mesoscale meteorology. *Atmospheric Chemistry and Physics*, 13(3), 1395–1410. <https://doi.org/10.5194/acp-13-1395-2013>
- Powley, H. R., Krom, M. D., & Van Cappellen, P. (2017). Understanding the unique biogeochemistry of the Mediterranean Sea: Insights from a coupled phosphorus and nitrogen model. *Global Biogeochemical Cycles*, 31, 1010–1031. <https://doi.org/10.1002/2017GB005648>
- Rengarajan, R., & Sarin, M. M. (2004). Atmospheric deposition fluxes of ⁷Be, ²¹⁰Pb and chemical species to the Arabian Sea and Bay of Bengal. *Indian Journal of Marine Sciences*, 33, 56–64.
- Ridame, C., Dekaezemacker, J., Guieu, C., Bonnet, S., L'Helguen, S., & Malien, F. (2014). Contrasted Saharan dust events in LNLC environments: Impact on nutrient dynamics and primary production. *Biogeosciences*, 11(17), 4783–4800. <https://doi.org/10.5194/bg-11-4783-2014>
- Rodríguez, S., Alastuey, A., Alonso-Pérez, S., Querol, X., Cuevas, E., Abreu-Afonso, J., et al. (2011). Transport of desert dust mixed with North African industrial pollutants in the subtropical Saharan air layer. *Atmospheric Chemistry and Physics*, 11(13), 6663–6685. <https://doi.org/10.5194/acp-11-6663-2011>
- Ruttenberg, K. C. (2003). The global phosphorus cycle. *Treatise on Geochemistry*, 8, 585–643. <https://doi.org/10.1016/B0-08-043751-6/08153-6>
- Sedwick, P. N., Sholkovitz, E. R., & Church, T. M. (2007). Impact of anthropogenic combustion emissions on the fractional solubility of aerosol iron: Evidence from the Sargasso Sea. *Geochemistry, Geophysics, Geosystems*, 8, Q10Q06. <https://doi.org/10.1029/2007GC001586>
- Shangguan, W., Dai, Y., Duan, Q., Liu, B., & Yuan, H. (2014). A global soil data set for Earth system modeling. *Journal of Advances in Modeling Earth Systems*, 6, 249–263. <https://doi.org/10.1002/2013ms000293>
- Sinha, P., Jaeglé, L., Hobbs, P. V., & Liang, Q. (2004). Transport of biomass burning emissions from southern Africa. *Journal of Geophysical Research*, 109, D20204. <https://doi.org/10.1029/2004JD005044>
- Slinn, W. G. N. (1982). Predictions for particle deposition to vegetative canopies. *Atmospheric Environment*, 16(7), 1785–1794. [https://doi.org/10.1016/0004-6981\(82\)90271-2](https://doi.org/10.1016/0004-6981(82)90271-2)
- Smits, M. M., Bonneville, S., Benning, L. G., Banwart, S. A., & Leake, J. R. (2012). Plant-driven weathering of apatite—The role of an ectomycorrhizal fungus. *Geobiology*, 10(5), 445–456. <https://doi.org/10.1111/j.1472-4669.2012.00331.x>
- Srinivas, B., & Sarin, M. M. (2015). Atmospheric deposition of phosphorus to the northern Indian Ocean. *Current Science*, 456–457, 104–114. <https://doi.org/10.1016/j.scitotenv.2013.03.068>
- Stibal, M., Anesio, A. M., Blues, C. J. D., & Tranter, M. (2009). Phosphatase activity and organic phosphorus turnover on a high Arctic glacier. *Biogeosciences*, 6(5), 913–922. <https://doi.org/10.5194/bg-6-913-2009>
- Stockdale, A., Krom, M. D., Mortimer, R. J., Benning, L. G., Carslaw, K. S., Herbert, R. J., et al. (2016). Understanding the nature of atmospheric acid processing of mineral dusts in supplying bioavailable phosphorus to the oceans. *Proceedings of the National Academy of Sciences of the United States of America*, 113(51), 14,639–14,644. <https://doi.org/10.1073/pnas.1608136113>
- Tamatamah, R. A., Hecky, R. E., & Duthie, H. (2005). The atmospheric deposition of phosphorus in Lake Victoria (East Africa). *Biogeochemistry*, 73(2), 325–344. <https://doi.org/10.1007/s10533-004-0196-9>
- Thingstad, T. F., Krom, M. D., Mantoura, R. F., Flaten, G. A., Groom, S., Herut, B., et al. (2005). Nature of phosphorus limitation in the ultraoligotrophic eastern Mediterranean. *Science*, 309(5737), 1068–1071. <https://doi.org/10.1126/science.1112632>
- Tyrrill, T. (1999). The relative influences of nitrogen and phosphorus on oceanic primary production. *Nature*, 400(6744), 525–531. <https://doi.org/10.1038/22941>
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., et al. (2010). Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009). *Atmospheric Chemistry and Physics*, 10(23), 11,707–11,735. <https://doi.org/10.5194/acp-10-11707-2010>
- Vergara-Temprado, J., Murray, B. J., Wilson, T. W., O'Sullivan, D., Browne, J., Pringle, K. J., et al. (2017). Contribution of feldspar and marine organic aerosols to global ice nucleating particle concentrations. *Atmospheric Chemistry and Physics*, 17(5), 3637–3658. <https://doi.org/10.5194/acp-17-3637-2017>
- Vet, R., Artz, R. S., Carou, S., Shaw, M., Ro, C.-U., Aas, W., et al. (2014). A global assessment of precipitation chemistry and deposition of sulfur, nitrogen, sea salt, base cations, organic acids, acidity and pH, and phosphorus. *Atmospheric Environment*, 93, 3–100. <https://doi.org/10.1016/j.atmosenv.2013.10.060>

- Wang, R., Balkanski, Y., Bopp, L., Aumont, O., Boucher, O., Ciais, P., et al. (2015). Influence of anthropogenic aerosol deposition on the relationship between oceanic productivity and warming. *Geophysical Research Letters*, *42*, 10,745–10,754. <https://doi.org/10.1002/2015GL066753>
- Yang, X., Post, W. M., Thornton, P. E., & Jain, A. (2013). The distribution of soil phosphorus for global biogeochemical modeling. *Biogeosciences*, *10*(4), 2525–2537. <https://doi.org/10.5194/bg-10-2525-2013>
- Yogev, T., Rahav, E., Bar-Zeev, E., Man-Aharonovich, D., Stambler, N., Kress, N., et al. (2011). Is dinitrogen fixation significant in the Levantine Basin, East Mediterranean Sea? *Environmental Microbiology*, *13*(4), 854–871. <https://doi.org/10.1111/j.1462-2920.2010.02402.x>
- Zamora, L. M., Prospero, J. M., Hansell, D. A., & Trapp, J. M. (2013). Atmospheric P deposition to the subtropical North Atlantic: Sources, properties, and relationship to N deposition. *Journal of Geophysical Research: Atmospheres*, *118*, 1546–1562. <https://doi.org/10.1002/jgrd.50187>
- Zhang, G., Zhang, J., & Liu, S. (2007). Characterization of nutrients in the atmospheric wet and dry deposition observed at the two monitoring sites over Yellow Sea and East China Sea. *Journal of Atmospheric Chemistry*, *57*(1), 41–57. <https://doi.org/10.1007/s10874-007-9060-3>
- Zhang, Y., Mahowald, N., Scanza, R. A., Journet, E., Desboeufs, K., Albani, S., et al. (2015). Modeling the global emission, transport and deposition of trace elements associated with mineral dust. *Biogeosciences*, *12*(19), 5771–5792. <https://doi.org/10.5194/bg-12-5771-2015>

Erratum

The caption of Figure 2(b) as originally published misstated the annual deposition rate of dust as $\text{g/m}^2/\text{year}^{-1}$. This error has since been corrected, and this may be considered the authoritative version of record.